

NONOPIOID TREATMENTS FOR CHRONIC PAIN

PRINCIPLES OF CHRONIC PAIN TREATMENT


Patients with pain should receive treatment that provides the greatest benefit. Opioids are not the first-line therapy for chronic pain outside of active cancer treatment, palliative care, and end-of-life care. Evidence suggests that nonopioid treatments, including nonopioid medications and nonpharmacological therapies can provide relief to those suffering from chronic pain, and are safer. Effective approaches to chronic pain should:

Use nonopioid therapies to the extent possible

Identify and address co-existing mental health conditions (e.g., depression, anxiety, PTSD)

Focus on functional goals and improvement, engaging patients actively in their pain management

Use disease-specific treatments when available (e.g., triptans for migraines, gabapentin/pregabalin/duloxetine for neuropathic pain)



Use first-line medication options preferentially

Consider interventional therapies (e.g., corticosteroid injections) in patients who fail standard non-invasive therapies

Use multimodal approaches, including interdisciplinary rehabilitation for patients who have failed standard treatments, have severe functional deficits, or psychosocial risk factors

NONOPIOID MEDICATIONS

MEDICATION	MAGNITUDE OF BENEFITS	HARMS	COMMENTS
Acetaminophen	Small	Hepatotoxic, particularly at higher doses	First-line analgesic, probably less effective than NSAIDs
NSAIDs	Small-moderate	Cardiac, GI, renal	First-line analgesic, COX-2 selective NSAIDs less GI toxicity
Gabapentin/pregabalin	Small-moderate	Sedation, dizziness, ataxia	First-line agent for neuropathic pain; pregabalin approved for fibromyalgia
Tricyclic antidepressants and serotonin/norepinephrine reuptake inhibitors	Small-moderate	TCAs have anticholinergic and cardiac toxicities; SNRIs safer and better tolerated	First-line for neuropathic pain; TCAs and SNRIs for fibromyalgia, TCAs for headaches
Topical agents (lidocaine, capsaicin, NSAIDs)	Small-moderate	Capsaicin initial flare/burning, irritation of mucus membranes	Consider as alternative first-line, thought to be safer than systemic medications. Lidocaine for neuropathic pain, topical NSAIDs for localized osteoarthritis, topical capsaicin for musculoskeletal and neuropathic pain

RECOMMENDED TREATMENTS FOR COMMON CHRONIC PAIN CONDITIONS

Low back pain

Self-care and education in all patients; advise patients to remain active and limit bedrest

Nonpharmacological treatments: Exercise, cognitive behavioral therapy, interdisciplinary rehabilitation

Medications

- First-line: acetaminophen, non-steroidal anti inflammatory drugs (NSAIDs)
- Second-line: Serotonin and norepinephrine reuptake inhibitors (SNRIs)/tricyclic antidepressants (TCAs)

Migraine

Preventive treatments

- Beta-blockers
- TCAs
- Antiseizure medications
- Calcium channel blockers
- Non-pharmacological treatments (Cognitive behavioral therapy, relaxation, biofeedback, exercise therapy)
- Avoid migraine triggers

Acute treatments

- Aspirin, acetaminophen, NSAIDs (may be combined with caffeine)
- Antinausea medication
- Triptans-migraine-specific

Neuropathic pain

Medications: TCAs, SNRIs, gabapentin/pregabalin, topical lidocaine

Osteoarthritis

Nonpharmacological treatments: Exercise, weight loss, patient education

Medications

- First-line: Acetaminophen, oral NSAIDs, topical NSAIDs
- Second-line: Intra-articular hyaluronic acid, capsaicin (limited number of intra-articular glucocorticoid injections if acetaminophen and NSAIDs insufficient)

Fibromyalgia

Patient education: Address diagnosis, treatment, and the patient's role in treatment

Nonpharmacological treatments: Low-impact aerobic exercise (e.g., brisk walking, swimming, water aerobics, or bicycling), cognitive behavioral therapy, biofeedback, interdisciplinary rehabilitation

Medications

- FDA-approved: Pregabalin, duloxetine, milnacipran
- Other options: TCAs, gabapentin

